

Age Adjusted Death Rates for Cerebrovascular Disease (Stroke) for Selected Population Groups, 1999-2001

| Indicator | White Males | White Females | Minority Males | Minority Females | Overall |
|-----------------------------|--------------------|----------------------|-----------------------|-------------------------|----------------|
| 99-01 Bladen Cnty Rate | 108.5 | 99.6 | 150.1 | 124.7 | 114.3 |
| 99-01 Brunswick Cnty Rate | 50.8 | 62.4 | 64 | 47.3 | 59.2 |
| 99-01 Columbus Cnty Rate | 88.1 | 78.1 | 89.1 | 61.1 | 77.8 |
| 99-01 Duplin Cnty Rate | 77.2 | 105.8 | 139.8 | 122.3 | 107.2 |
| 99-01 Harnett Cnty Rate | 67.3 | 62.3 | 116.3 | 52.2 | 67.7 |
| 99-01 New Hanover Cnty Rate | 52.4 | 75.7 | 124.3 | 66 | 72.2 |
| 99-01 Onslow Cnty Rate | 68.3 | 59.5 | 74.9 | 79.6 | 66.6 |
| 99-01 Pender Cnty Rate | 109 | 86.5 | 126.3 | 101.5 | 95.7 |
| 99-01 Robeson Cnty Rate | 120.7 | 95.1 | 105.2 | 90.5 | 101.4 |
| 99-01 Sampson Cnty Rate | 91.4 | 79 | 111.1 | 104.5 | 91.3 |
| 99-01 Horry Cnty Rate | 63 | 70.1 | 112.6 | 116.8 | 75.6 |
| 99-01 NC Rate | 69.3 | 67.9 | 103.9 | 82.8 | 73.6 |
| 99-01 SC Rate | 72.3 | 70.2 | 126.5 | 101.7 | 80.9 |

